

SUPPLEMENTAL PRODUCT BULLETIN



Biochemical Repellent for Sciarid and Phorid Fly Control

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

EPA Exempted Product / FIFRA 25(b)

Natural Forces, LLC
PO Box 2601
Davidson NC 28036-2601

Please refer to the Divert-O™ label for First Aid, precautionary statements, and environmental hazards.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of the application. Read the label affixed to the container for Divert-O™ before applying. Use of Divert-O™ according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Divert-O™.

GENERAL INFORMATION

Divert-O repels mushroom flies (Sciarid and Phorid adults) in mushroom houses and adjacent premise areas. Use prior to filling the house, during filling, during cool-down, during spawning, up to pinning, during and between breaks. Treat when flies appear or conditions are favorable for attracting flies. Maintain application during entire period of time necessary to repel flies. Flies will re-enter areas immediately after fragrance subsides.

SURFACE APPLICATION

Spray outside and inside walls, floors, openings, HVAC intakes and sideboards of mushroom houses. Also spray over the plastic covering the beds and trays after spawning.

Hand Application Directions (One Gallon): To obtain maximum benefits apply every seven (7) to ten (10) days before infestation begins or as needed. If infestation is present, apply 0.5 - 4.0 fl. oz. of Divert-O per gallon of water with emulsifier. Apply as a full coverage, making sure all surfaces are covered. Apply as a preventative treatment prior to insect infestation.

Must be mixed with emulsifier for uniform dispersal.

High Volume Machine and Hose End Sprayer: Apply by ground machine at a rate of 1.5 quarts to 2 gallons of Divert-O per 100 gallons. Apply as a full coverage, making sure both sides of the surfaces are covered. Apply as a preventative treatment prior to insect infestation.

Must be mixed with emulsifier for uniform dispersal.

Application: Do not apply this product through any type of irrigation system.